

Pharmaceutical classification	Over the counter drug
Medicinal effect classification	Antipyretic Analgesics
product name	バイエルアスピリン BAYER ASPIRIN
Product features	Bayer Aspirin is a non-pyrine antipyretic analgesic developed by the German company Bayer. The active ingredient aspirin (acetylsalicylic acid) suppresses the generation of substances that cause pain and fever. The aspirin in Bayer Aspirin is made from small, uniform crystals. These dissolve quickly in the gastrointestinal tract for fast absorption, making it very effective against pain and fever.
Indication	(1) Relief of headache, toothache, pain after tooth extraction, sore throat (throat pain), earache, joint pain, neuralgia, lumbago, muscular pain, pain due to stiff shoulders, contusion pain, bone fracture pain, pain associated with sprain (sprain pain), painful menses (menstrual pain), and traumatic pain (2) Relief of fever at the time of chills (feeling cold due to fever) and fever
Dosage and administration	When symptoms appear, take the following single dose. Avoid taking on an empty stomach as much as possible. Always leave at least 4 hours between doses.  < Age: single dose: number of daily doses > Adults (over 15): 1 tablet: 3x as the limit Under 15 years old: Do not take  ★Take with a glass of water.
Precautions of Dosage and Administration	(1)Strictly follow the prescribed dosage and administration. (2) How to remove tablets from the package Push the convex part of the PTP sheet containing the tablets with your fingertips to break the aluminum foil on the back, remove it and take. (If accidentally swallowed without removing the packag, it will lead to unexpected accidents such as piercing the esophageal mucosa.)
Ingredient and amount	Bayer Aspirin is a white tablet containing the following ingredients. Aspirin : 500mg : Reduces fever and alleviates pain.
Precautions	<p>■When not to use the product (If you do not follow these instructions, the current symptoms may worsen or adverse reactions/incidents are more likely to occur.)</p> <p>1. This product should not be taken by the following persons. (1)Those who have experienced allergic symptoms (rash, redness, itching, edema, etc.) to this medication or any of its ingredients (2)Patients who have experienced asthma from taking this drug, other antipyretic analgesics, cold remedies. (3)Children under 15 years old. (4)Those with gastric or duodenal ulcers. (5)Those who tend to bleed easily (spotting on hands and legs, tendency toward purpura, etc.). (6)Pregnant women who are expected to given birth within 12 weeks.</p> <p>2. This drug should not be taken together with the following drugs: Other antipyretics/analgesics, cold medicine, sedatives</p> <p>3. Do not drink alcohol before/after taking this medicine</p> <p>4. Do not take this medicine for a long time</p> <p>■Consultation</p> <p>1. The following persons should contact a physician, dentist, pharmacist, or registered salesperson for a consultation before administration. (1)Patients undergoing medical treatment from a physician or dentist. (2)Pregnant women or women suspected of being pregnant. (3)Nursing women. (4)The elderly (5)Patients who have experienced allergic symptoms associated with drugs, etc. (6)Persons diagnosed as having the following heart disease, kidney disease, liver disease (7)Patients who experienced any of the following diseases. gastric/duodenal ulcer</p> <p>2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document to your physician, pharmacist, or registered salesperson for a consultation.</p>

	<p>&lt;Related parts : Symptoms&gt;</p> <p>Skin : rash/redness,itching, bruised</p> <p>Gastrointestinal system : nausea and vomiting, loss of appetite, heartburn, stomach heaviness, stomachache,abdominal pain, diarrhea, bloody stool, gastrointestinal bleeding</p> <p>Neuropsychiatric system : dizziness</p> <p>Other : nose bleeds, bleeding of the gums, difficult to stop bleeding, bleeding, fever, sore throat, back pain, excessive temperature decrease, edema, anemia, ear ringing, hearing loss</p> <p>The following serious symptoms may occur in rare cases. In such cases, immediately seek medical aid:</p> <p>&lt;Symptom name : Symptoms&gt;</p> <p>Shock (anaphylaxis): Symptoms, such as itching of skin, urticaria, hoarseness, sneezing, itchy throat, breathing difficulties, palpitations, and clouding of consciousness may occur immediately after take.</p> <p>Oculomucocutaneous syndrome (Stevens-Johnson syndrome) , toxic epidermal necrolysis: hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, etc. may persist or suddenly worsen.</p> <p>Acute coronary syndrome associated with allergic reactions: Symptoms, such as constricting chest pain, shortness of breath, a sensation of strong pressure on the chest, jaw pain, left arm pain, chills, dizziness, sweating, fever, decreased level of consciousness, swelling around the lips, itching, hives, rash, itching of the throat, palpitations, and others.</p> <p>Hepatic function failure: Symptoms, such as fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, general malaise, loss of appetite, etc. may occur.</p> <p>Asthma: Symptoms, such as wheezing or hissing when breathing, and difficult breathing, etc. may occur.</p> <p>Aplastic anemia: Symptoms, such as bruises, nose bleeds, bleeding gums, fever, pale appearance of skin and mucosa, fatigue, palpitations, shortness of breath, becoming dizzy from feeling sick and blood in the urine, etc. may occur.</p> <p>3. When symptoms do not improve even after taking the medicine 5 to 6 times, stop taking this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this instruction leaflet with you.</p>
Additive	Cellulose, corn starch
Precautions for storage and handling	<p>(1)Store in a cool, low humidity place away from direct sunlight.</p> <p>(2)Keep out of reach of children.</p> <p>(3)Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug. )</p> <p>(4)Do not take the product after its expiration date.</p>
Other	
Manufacturing and sales company	Bayer YakuhinCo., Ltd. 2-4-9 Moto Umeda, Kita-ku, Osaka
Sales company	Sato Pharmaceutical Co., Ltd. 1-5-27 Moto Akasaka, Minato-ku, Tokyo
Risk category	Designated Category 2 OTC medicine
Disclaimer on Multilingual Product Information	<ul style="list-style-type: none"> <li>• This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.</li> <li>• Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.</li> <li>• Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.</li> <li>• Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.</li> </ul>