

Before using this medicine, be sure to read this written instruction.
Also please keep it in a safe place so that you can read it when necessary.

Common cold medicine

Designated Class 2 OTC Medicine

For sore throat, fever, runny nose, nasal congestion, cough, and phlegm

ルルAプレミアム

Lulu® A Premium

Features of Lulu A Premium

- This product is a comprehensive cold medicine that is highly effective against all 11 cold symptoms. It may be taken by adults and children 7 years of age and over.
- Tranexamic acid, an anti-inflammatory ingredient, and acetaminophen, an antipyretic analgesic, provide excellent relief for sore throat and fever.
- Clemastine fumarate, a long-acting antihistamine ingredient, and belladonna total alkaloids, a parasympathetic blocker, provide dual relief for runny nose and nasal congestion through their combined action.
- Bromhexine hydrochloride, an expectorant ingredient, helps expel phlegm, one of the causes of coughs.
- Dextromethorphan hydrobromide hydrate, an antitussive ingredient, provides excellent relief for persistent coughs.
- These sugar-coated tablets have no bitter taste and are easy to swallow even for those who do not like taking medicine.
- This product comes in convenient PTP packaging that is easy to carry when going out or on the go.



Precautions



Things to avoid

(If you fail to follow these precautions, your present symptoms may worsen, or side effects/accidents may occur more readily.)

1. Do not take this medicine if:

- (1) You have experienced allergic reactions to this medicine or its components in the past
- (2) You have developed asthma after taking this or other cold medicines, or antipyretic analgesics.

2. Do not use any of the following medicines while taking this medicine:

Other cold medicines, antipyretic analgesics, sedatives, antitussives, expectorants, oral medicines containing antihistamine (oral medicines for rhinitis, motion sickness medicines, anti-allergic medicines, etc.), gastrointestinal analgesic/ antispasmodic medicines, oral medicines containing tranexamic acid

3. Do not drive a vehicle or operate machinery after taking this medicine

(Symptoms such as sleepiness, blurred vision, or abnormal dazzling may occur.)

4. Do not consume alcohol before or after taking this medicine.

5. Do not use this product continuously for a long term.

* 6. Do not take more than the specified dose.



Things to consult

1. Consult with a doctor, pharmacist, or registered salesclerk before taking this medicine if:

- (1) You are receiving medical or dental treatment.
- (2) You are pregnant or might be pregnant.

(3) Breastfeeding women.

(4) You are elderly.

(5) You have experienced allergic reactions to medications before.

(6) You are experiencing any of the following symptoms: High fever, dysuria

(7) You have been diagnosed with any of the following conditions:

Thyroid functional disorder, diabetes, heart disease, hypertension, liver disease, kidney disease, Gastric/duodenal ulcer, glaucoma, thrombosis (cerebral thrombosis, myocardial infarction, thrombophlebitis, etc.) or you are at risk of developing thrombosis

2. If any of the following symptoms occur after taking this medicine, they may indicate side effects. Discontinue use immediately and consult with a doctor, pharmacist, or registered salesclerk with this instruction leaflet.

Related Part	Symptoms
Skin	Rash/redness, itching
Digestive system	Nausea/vomiting, heartburn, loss of appetite
Neuropsychiatric system	Dizziness, excitement, convulsions, headache
Respiratory system	Shortness of breath, difficulty breathing
Urinary system	Dysuria
Other	Excessive temperature decrease, facial flushing, abnormal sensitivity to light

In rare cases, the following severe symptoms may occur. In such cases, seek medical attention immediately.

Name of Symptom	Symptoms
Shock (anaphylaxis)	Symptoms such as itchy skin, urticaria, hoarseness, sneezing, itchy throat, difficulty breathing, palpitations, and clouding of consciousness may occur immediately after taking this medicine.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), Toxic epidermal necrolysis, Acute generalized exanthematous pustulosis	Hyperthermia, ocular hyperaemia, eye discharge, lip erosion, pain throat, widespread skin rash/redness, small pimples (small pustules) on reddened skin, general malaise, anorexia, etc. may persist or suddenly worsen.
Drug-Induced Hypersensitivity Syndrome (DIHS)	Widespread skin redness, systemic rash, fever, general malaise, swelling of lymph nodes (neck, armpits, groin, etc.)
Liver dysfunction	Symptoms such as fever, itching, rash, jaundice (skin and whites of the eyes turn yellow), brown urine, general malaise, loss of appetite, etc. may occur.
Kidney disorders	Symptoms such as fever, rash, reduced urinary output, full-body swelling, general malaise, joint pain and diarrhea, etc. may occur.
Interstitial pneumonia	Shortness of breath or difficulty breathing when climbing stairs or straining yourself a little, sudden or persistent symptoms like dry cough, fever, etc.
Asthma	Wheezing or hissing when breathing, and difficult breathing, etc. may occur.

3. The following symptoms may occur after taking this medicine. If these symptoms persist or worsen, discontinue use and bring these instructions to a consultation with your doctor, pharmacist, or registered salesclerk.

Diarrhea, constipation, dry mouth, drowsiness, blurred vision

4. If symptoms do not improve after 5-6 doses, stop taking this medicine immediately and bring these instructions to a consultation with your doctor, pharmacist, or registered salesclerk.

Indications

Relief of various cold symptoms (sore throat, fever, runny nose, nasal congestion, cough, phlegm, chills, headache, sneezing, joint pain, muscle pain)

Dosage and Administration

Take the following amount with water or warm water.

Age	Amount per dose	Number of doses per day
Adults (15 years or older)	3 tablets 	3 times Take within 30 minutes after a meal if possible.
11 to 14 years	2 tablets 	
7 to 10 years	1 tablets 	
Under 7 years	Do not take.	

<Precautions related to Dosage and Administration>

- (1) Strictly follow the Dosage and Administration.
- (2) Use by children of 7 years and over should be supervised by a parent or guardian.
- (3) How to take out the tablets:

As shown in the figure on the right, use your finger to push the PTP sheet that contains the tablet. Take off the aluminum foil on the back and ingest the tablet. (If you mistakenly take the tablet with the wrapping on it, it could puncture the mucous membrane of the esophagus and lead to other unknown consequences.)



Ingredients and Quantities

This product consists of white sugar-coated tablets containing the following ingredients per 9 tablets.

Ingredient	Quantity	Function
Clemastine Fumarate	1.34 mg	Suppresses allergic cold symptoms for an extended period (runny nose, sneezing, etc.).
Belladonna Total Alkaloids	0.6 mg	Suppresses runny nose.
Bromhexine Hydrochloride	12 mg	Helps expel phlegm, one of the causes of coughs.
Tranexamic Acid	750 mg	Suppresses [plasmin], a source of inflammation and soothes swelling and pain in the throat.
Acetaminophen	900 mg	Reduces fever and relieves sore throat, headache, and joint pain.
dl-Methylephedrine Hydrochloride	60 mg	Expands the bronchial tubes, relieves coughs.
Dextromethorphan Hydrobromide Hydrate	48 mg	Works on the cough center to relieve coughs.

Additives: Cellulose, Hydroxypropylcellulose, Magnesium Aluminometasilicate, Carmellose Calcium, Magnesium Stearate, Fumaric Acid, Polyvinylacetal Diethylaminoacetate, Stearic Acid, Hypromellose, White Soft Sugar, Talc, Partially Hydrolyzed Polyvinyl Alcohol, Povidone, Acacia, Titanium Oxide, Carnauba Wax

Storage and Handling Precautions


- (1) Store in a cool and dry place and out of direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer and keep in a different container. (It can lead to misuse or lowering of quality.)
- (4) Please do not use the product after the indicated expiration date. Also, after opening the foil pouch, use the product within 12 months.
- (5) Write the opening date in the "Date Opened" field on the box.

Common Colds Causes

A cold is an acute inflammatory disease of the upper respiratory tract, including the throat and nose.

Viruses and other substances enter the body through the throat or nose, causing an excessive inflammatory response. This inflammatory response can cause respiratory symptoms such as sore throat, runny nose, cough, and phlegm, as well as fever.

Firstly, controlling and calming inflammation is the key to relieving cold symptoms.



Image

If you have any questions about this product, please contact the store where you purchased it or at the following address:

Daiichi Sankyo Healthcare Co., Ltd. Customer Service Center

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan zip/postal code 103-8234

TEL: 0120-337-336

Reception Hours: 9:00 am - 5:00 pm (closed on Saturdays, Sundays, and holidays)

For details about Lulu, visit www.lulu.jp

02® Registered Trademark



Manufacturer & Marketing Authorization Holder

Daiichi Sankyo Healthcare Co., Ltd.

3-14-10, Nihonbashi, Chuo-ku, Tokyo, Japan

Homepage: <https://www.daiichisankyo-hc.co.jp/>