

Make sure to read this instruction sheet before use.
Additionally, store this sheet so that it can be referenced when needed.



Therapeutic agent for recurrent vaginal candidiasis

MEDITREAT

Class 1 OTC drug

<Externally applied medicine for vagina>

- ◆ MEDITREAT is a medicine for treating recurrent vaginal candidiasis, which is mainly composed of miconazole nitrate.
- ◆ Miconazole nitrate is effective against Candida, which causes vaginal candidiasis, and is used to treat vaginal candidiasis.
- ◆ Effective against itchiness in and discharge from sensitive areas caused by recurrent vaginal candidiasis.



Precautions for Use



Things You Must Not Do



(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects occurring)

1 The following individuals should not use this product:

- (1) Persons who think this is their first onset of symptoms. (The first onset of symptoms may be caused by other conditions, and a diagnosis from a doctor is required in such cases)
- (2) Persons who have had allergic symptoms as a result of using this product or any of its components. (Allergic symptoms may occur again as a result of using this product)
- (3) Persons under 15 years or over 60 years of age. (For persons under 15 years of age, the symptoms are highly likely to be their first onset. For persons over 60 years of age, the symptoms may be caused by other conditions or mixed infection due to other bacteria)
- (4) Persons who are pregnant or who think they may be pregnant. (Medication use should be considered carefully and a diagnosis from a doctor is required)
- (5) Persons with fever, chills, abdominal pain, mid- or upper back pain, colored, bloody, or fishy-smelling vaginal discharge, absence of periods, irregular or abnormal vaginal bleeding, or ulcers, swelling or inflammation in the vagina or vulvar area. (The symptoms may be caused by other conditions, and a diagnosis from a doctor is required)
- (6) Persons who have received the following diagnoses:
Diabetes (The infection is likely to recur frequently, and a diagnosis from a doctor is required)
- (7) Persons taking warfarin or other anticoagulants. (The effects of warfarin may increase the tendency to bleed)
- (8) Persons who get this infection frequently. (Once every 1 to 2 months or at least twice in 6 months)
- (9) Persons who are uncertain whether the symptoms are caused by recurrent vaginal candidiasis. (If it is difficult to determine yourself, a diagnosis from a doctor is required)

2 Do not use on the following areas:

- (1) Anywhere other than inside the vagina. (This product is only effective against vaginal candidiasis caused by Candida)

3 Do not use the following medicines while using this product:

- (1) Any dermatological agent other than therapeutic agents for candidiasis. (Your current symptoms may worsen or your treatment may be delayed)



Things To Consult On

1 The following persons should consult a doctor or pharmacist before use:

- (1) Persons receiving treatment from a doctor. (The medication prescribed by your doctor may be affected or you may be using the same medication as this product)
- (2) Persons with a history of allergic reactions caused by medicines, etc. (This product may cause allergic reactions to persons with a history of allergic reactions caused by medicines, etc.)
- (3) Persons who are breastfeeding. (Medication use should be considered carefully)

2 If you experience any of the following symptoms after use, you may be experiencing side effects. Discontinue use immediately and consult a doctor or pharmacist with a copy of this instruction sheet.

Relevant Area	Symptoms
Other than vagina	Hives, itching

3 The following symptoms may appear after using this product. If these symptoms persist or worsen, discontinue use and consult with your doctor or pharmacist, bringing this instruction sheet with you

Relevant Area	Symptoms
Vagina	Itching, redness, pain, burning sensation, irritation

(These symptoms may be allergic reactions to this product or the pharmacological action of this product may have developed strongly, and continued use of the same medicine may further worsen symptoms in such cases)

4 If you do not notice improvement in your symptoms after 3 days of use or your symptoms remain after 6 days of use, consult your doctor. (Your symptoms may be severe or may be caused by other conditions)

Efficacy

Recurrent vaginal candidiasis (limited to persons who have been diagnosed and treated by a doctor in the past)

[Explanation]

Vaginal candidiasis is a fungal infection in the vaginal area caused by Candida yeast, which causes itching of the vagina, white vaginal discharge, redness, a burning sensation and pain.

Usage/Dosage

Adults (15 years or older and under 60 years of age): Insert 1 suppository deep inside the vagina once daily before bed. Use the product continuously for 6 days.

However, if you do not notice improvement in your symptoms after 3 days of use or your symptoms remain after 6 days of use, consult your doctor.

*Bathe or clean the affected area with warm water before use.


Age	Single dose	Number of uses
15 years or older and under 60 years of age	 1	Once per day (Before bed)
Under 15 years or over 60 years of age	Do not use this product	

Note:


- (1) Strictly follow the prescribed usage and dosage.
- (2) If the suppository is soft, allow it to cool for a while before use. If it is too hard, wait until it is softer before use.
- (3) This product should only be used inside the vagina.
- (4) Wash hands thoroughly before and after use.
- (5) Do not reuse the applicator. (Reusing the applicator is unpreferable from a hygiene standpoint)
- (6) Even if your symptoms start to resolve, use the product for 6 days from the start of use.
- (7) Do not use the product during your period. If your period starts during use, discontinue use. In such cases, consult a doctor to confirm whether the infection has been treated. (The medicine may flow out during a period, reducing its effects)

How to insert and remove a suppository


1. Tear off 1 piece along the dotted line.



2. Take out the suppository from the container by pulling the sides apart.



3. Insert the suppository into the vagina.



Ingredients

In 1 suppository

Ingredients	Amount	Action
Miconazole nitrate	100 mg	Highly effective against Candida.

Additives: Hard fat

Precautions for Storage and Handling

- (1) Store in a cool, dry place at 30°C or below, out of direct sunlight. (This product melts at body temperature. Store it in a refrigerator if possible after purchase)
- (2) Store this product out of reach of children.
- (3) When storing, place the tip of the suppository facing downward in the package, and store the package vertically according to the marking on the box. (To prevent deformation of suppository)
- (4) Avoid contact with latex contraceptive products, such as condoms and pessaries. (These products may be damaged or deteriorate.)
- (5) Do not use this product beyond its expiration date. (In order to ensure quality.)



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.