



Make sure to read this instruction sheet before taking this medicine. Additionally, store this sheet so that it can be referenced when needed.

NARON ACE

PREMIUM

Also works on recurring headaches

Brand name: NARON ACE PREMIUM

Class **2** OTC drug <Antipyretic analgesic>

- ◆ NARON ACE PREMIUM is an antipyretic analgesic that contains 2 fever and pain-relieving ingredients combined with an ingredient that protects the stomach lining.
- ◆ With the combination of ibuprofen to block the source of pain and the largest amount (Note) of 1500 mg of etheznamide to stop the transmission of pain, NARON ACE PREMIUM also works against recurring painful headaches. (Note: In terms of daily amount in the NARON series)
- ◆ Dried aluminum hydroxide gel protects the stomach lining and prevents stomach irritation.



Precautions for Use

Things You Must Not Do



(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

- 1** The following persons should not take this medicine:
 - (1) Persons who have had allergic symptoms to this medicine or any of its ingredients.
 - (2) Persons who have experienced asthma while taking this medicine or other antipyretic analgesics or cold medicines.
 - (3) Children under 15 years of age.
 - (4) Pregnant women whose due date is within 12 weeks.
- 2** Do not take any of the following medicines while taking this medicine: Other antipyretic analgesics, cold medicines, sedatives
- 3** Do not drink alcohol before or after taking this medicine
- 4** Do not use for long periods of time



Things To Consult On



- 1** The following individuals should consult with their doctor, dentist, pharmacist or a registered seller before taking this medicine:
 - (1) Persons receiving treatment from a doctor or dentist.
 - (2) Persons who are pregnant or who think they may be pregnant.
 - (3) Persons who are breastfeeding.
 - (4) Elderly persons.
 - (5) Persons who have had allergic symptoms due to medication or similar substances in the past.
 - (6) Persons who have received any of the following diagnoses: Heart disease, kidney disease, liver disease, systemic lupus erythematosus, mixed connective tissue disease
 - (7) Persons who have had the following diseases: Gastric/duodenal ulcer, ulcerative colitis, Crohn's disease
- 2** If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult with your doctor, dentist, pharmacist or registered seller, bringing this instruction sheet with you

Relevant Area	Symptoms
Skin	Rash, redness, itching, bruising
Digestive system	Nausea/vomiting, loss of appetite, stomach discomfort, stomach pain, canker sores, heartburn, upset stomach, gastrointestinal bleeding, abdominal pain, diarrhea, bloody stool
Neuropsychiatric system	Dizziness
Circulatory system	Palpitations
Respiratory system	Shortness of breath
Other	Blurred vision, ringing in ears, swelling, nosebleed, bleeding gums, bleeding that does not stop easily, bleeding, back pain, excessively low body temperature, feeling sluggish

In rare cases, serious side effects listed on the right may occur. In such cases, see a doctor immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion immediately after taking this medicine.
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis	High fever, bloodshot eyes, eye discharge, lip sores, sore throat, widespread skin rash/redness, etc., that persists or worsens rapidly.
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general sluggishness, loss of appetite, etc.
Kidney damage	Fever, rash, decreased urine output, general swelling, general sluggishness, joint pain, diarrhea, etc.
Aseptic meningitis	Severe headache accompanied by stiffness in the neck, fever, nausea/vomiting, etc. (Such symptoms are particularly frequently reported in those receiving treatment for systemic lupus erythematosus or mixed connective tissue disease.)
Myocardial infarction*	Tightening of the chest, difficulty breathing and cold sweat.
Cerebrovascular disorder*	Sudden reduction or loss of consciousness, difficulty moving a hand or foot on one side, headache, vomiting, dizziness, difficulty speaking or forming words, etc.
Asthma	Difficulty breathing or a wheezing or whistling sound when breathing.
Aplastic anemia	Bruising, nosebleeds, bleeding gums, fever, pale skin and mucous membranes, fatigue, palpitations, shortness of breath, feeling unwell and dizzy, blood in urine, etc.
Agranulocytosis	Sudden high fever, chills, sore throat, etc.

3 The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, discontinue use and consult with your doctor, pharmacist, or registered seller, bringing a copy of this instruction sheet with you
Constipation

4 If symptoms do not improve after using this medicine 5 to 6 times, discontinue use and consult with your doctor, dentist, pharmacist or registered seller, bringing this instruction sheet with you

Efficacy

- ◆ Relief from headache, painful menses (menstrual pain), toothache, pain after tooth extraction, lower back pain, pain due to stiff shoulders, muscle pain, joint pain, pain associated with bruising, pain associated with sprains, pain associated with bone fractures, traumatic pain, nerve pain, sore throat (throat pain) and ear ache
- ◆ Fever relief during chills (feeling cold due to fever) and fevers



Usage/Dosage

Take the following dosage with room-temperature or lukewarm water. Avoid taking on an empty stomach as much as possible. Wait at least 4 hours between doses.

Age	Single dose	Number of doses
Adults (15 years and older)	2 tablets	Up to 3 times a day
Under 15 years old	Do not take this medicine	

Note:

- (1) Strictly follow the prescribed usage and dosage.
- (2) How to take out the tablets:
As shown in the illustration, press the convex part of the PTP sheet containing the tablet hard with a fingertip to tear the aluminum foil on the reverse side, remove the tablet, and take it. (Accidental ingestion of the PTP sheet as-is may lead to unexpected accidents, such as penetration into the esophageal mucosa.)



The medicine should be taken as follows:

Take as soon as the pain starts.



Take 2 tablets each time with a cup of room-temperature or lukewarm water.



As much as possible, eat or drink beforehand to avoid taking the medicine on an empty stomach.



Ingredients

In 2 tablets

Ingredients	Amount	Action
Ibuprofen	150 mg	The combination of these 2 ingredients relieves various types of pain and fever symptoms.
Ethenzamide	500 mg	
Dried aluminum hydroxide gel	66.7 mg (200 mg in 6 tablets)	Protects the stomach lining and prevents stomach irritation.

Additives: Cellulose, anhydrous silicic acid, magnesium aluminometasilicate, croscarmellose sodium, sodium starch glycolate, hypromellose, talc, hydroxypropyl cellulose, magnesium stearate

Precautions for Storage and Handling

- (1) Store in a cool, dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not replace the medicine into a different container. (This may result in misuse or a change in quality)
- (4) Do not take the product after the expiration date. Use this medicine within 6 months of opening, even if still within the expiration date. (In order to ensure quality)



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.