

**Make sure to read this instruction sheet before taking this medicine.
Additionally, store this sheet so that it can be referenced when needed.**

For severe throat pain, fever, cough and runny nose

PABRON LX TABLETS

Brand name: PABRON LX TABLETS

Pharmaceuticals
requiring guidance

<Cold medicine>

◆ PABRON LX TABLETS is a cold medicine containing 4 active ingredients including loxoprofen sodium hydrate and L-carbocysteine. It is effective against 11 symptoms of a bad cold, including sore throat and fever.



Precautions for Use



Things You Must Not Do



(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

1 The following persons should not take this medicine:

- (1) Persons who have had allergic symptoms to this medicine or any of its ingredients.
- (2) Persons who have experienced asthma while taking this medicine or other cold medicines or antipyretic analgesics.
- (3) Children under 15 years of age.
- (4) Persons being treated for the following at a medical facility:
Gastric/duodenal ulcer, liver disease, kidney disease, heart disease
- (5) Persons who have been told by a doctor that they have a blood abnormality (blood disease), such as a low number of red blood cells (anemia), a low number of platelets (bleeding does not stop easily or you are prone to bleeding) or a low number of white blood cells.

(6) Pregnant women whose due date is within 12 weeks.

2 Do not use any of the following medicines while taking this medicine:

Other cold medicines, antipyretic analgesics, sedatives, antitussive expectorants, oral medicines containing antihistamines, etc. (oral medicines for rhinitis, motion sickness medicines, allergy medicines, etc.)

3 Do not drive a vehicle or operate machinery after taking this medicine

(You may experience drowsiness, etc.)

4 Do not drink alcohol before or after taking this medicine

5 Do not take this medicine for more than 5 days



Things To Consult On

1 The following persons should consult with their doctor or pharmacist before taking this medicine:

- (1) Persons receiving treatment from a doctor or dentist.
- (2) Persons who are pregnant or who think they may be pregnant.
- (3) Persons who are breastfeeding.
- (4) Elderly persons.
- (5) Persons who have had allergic symptoms due to medication or similar substances in the past.
- (6) Persons with the following symptoms:
High fever, difficulty urinating
- (7) Persons who have received any of the following diagnoses:
Liver disease, glaucoma, bronchial asthma, ulcerative colitis, Crohn's disease, systemic lupus erythematosus, mixed connective tissue disease
- (8) Persons who have had the following diseases:
Gastric/duodenal ulcer, liver disease, kidney disease, blood disease

2 If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult a doctor or pharmacist, bringing this instruction sheet with you

- (1) If symptoms such as an excessive drop in body temperature, weakness (lack of strength) or cold extremities (cold hands or feet) occur after taking this medicine.
- (2) If you develop a peptic ulcer or swelling after taking this medicine. In addition, in rare cases, serious symptoms may occur, such as gastrointestinal bleeding (causing vomiting of blood, nausea/vomiting, abdominal pain, black tar-like stool, bloody stool, etc.), gastrointestinal perforation (a hole forms in the gastrointestinal tract causing nausea/vomiting, severe abdominal pain, etc.) or narrowing or obstruction of the small or large intestine (causing nausea/vomiting, abdominal pain, abdominal bloating, etc.). In such cases, see a doctor immediately.
- (3) If the following symptoms appear after taking this medicine:

In rare cases, the following serious symptoms may occur.
In such cases, see a doctor immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion immediately after taking this medicine.
Mucocutaneous ocular syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis, erythema multiforme, acute generalized exanthematous pustulosis	High fever, bloodshot eyes, eye discharge, lip sores, sore throat, widespread skin rash/redness, blisters appearing on red areas of the skin, small bumps (small pustules) appearing on reddened skin, whole body sluggishness and loss of appetite, etc., that persists or worsens rapidly.
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general sluggishness, loss of appetite, etc.
Kidney damage	Fever, rash, decreased urine output, general swelling, general sluggishness, joint pain, diarrhea, etc.
Aseptic meningitis	Severe headache accompanied by stiffness in the neck, fever, nausea/vomiting, etc. (Such symptoms are particularly frequently reported in those receiving treatment for systemic lupus erythematosus or mixed connective tissue disease.)
Interstitial pneumonia	Shortness of breath or difficulty breathing on exertion, such as when climbing stairs, dry cough and fever, etc. These symptoms may appear suddenly or persist.
Asthma	Difficulty breathing or a wheezing or whistling sound when breathing.
Blood disorder	Sore throat, fever, general sluggishness, pale face and whitish inner eyelids, becoming prone to bleeding (bleeding gums, nosebleeds, etc.), and bruising (the color does not disappear even when pressed on), etc.
Congestive heart failure	General sluggishness, palpitations, shortness of breath, chest discomfort, chest pain, dizziness and fainting, etc.
Myocardial infarction*	Tightening of the chest, difficulty breathing and cold sweat.
Cerebrovascular disorder*	Sudden reduction or loss of consciousness, difficulty moving a hand or foot on one side, headache, vomiting, dizziness, difficulty speaking or forming words, etc.

Relevant Area	Symptoms
Skin	Rash, redness, itching
Digestive system	Nausea/vomiting, loss of appetite, abdominal pain, stomach discomfort, abdominal bloating, heartburn, canker sores, indigestion
Neuropsychiatric system	Dizziness, numbness, headache
Circulatory system	Increased blood pressure, palpitations
Urinary tract	Difficulty urinating, blood in urine
Other	Chest pain, fatigue, facial flushing, fever, anemia

Rhabdomyolysis	Muscle pain in the limbs, shoulders and lower back, etc., numbness in the limbs, weakness, stiffness, general sluggishness, reddish-brown urine, etc.
----------------	---

3 The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, discontinue use and consult with your doctor or pharmacist, bringing this instruction sheet with you

Dry mouth, constipation, diarrhea, drowsiness

4 If symptoms do not improve after taking this medicine 5 to 6 times, discontinue use and consult with your doctor or pharmacist, bringing this instruction sheet with you (especially if fever lasts for more than 3 days or occurs repeatedly)

Usage/Dosage

Take the following dosages with room-temperature or lukewarm water within no more than 30 minutes after meals.

Age	Single dose	Number of doses
15 years and older	2 tablets 	3 times a day
Under 15 years old	Do not take this medicine	

Note:

(1) Strictly follow the prescribed usage and dosage.

(2) How to take out the tablets:

As shown in the illustration, press the convex part of the PTP sheet containing the tablet hard with a fingertip to tear the aluminum foil on the reverse side, remove the tablet, and take it. (Accidental ingestion of the PTP sheet as-is may lead to unexpected accidents, such as penetration into the esophageal mucosa.)

<Tablet removal diagram>
Push it out



Efficacy

Relieves cold symptoms (sore throat, fever, cough, phlegm, runny nose, stuffy nose, sneezing, chills, headache, joint pain, muscle pain)

For sore throat, fever, etc.



For cough/phlegm



For sneezing, runny nose, stuffy nose



Ingredients

In 2 tablets

Loxoprofen sodium hydrate	68.1 mg (60 mg as anhydrous)	Relieves fever and pain, including fever, headache and sore throat.
L-carbocysteine	250 mg	Returns airway mucus and mucous membranes toward their normal state.
Tipecidine hibenazate	25 mg	Acts on the cough center in the brain to relieve cough.
Mequitazine	1.33 mg (4 mg in 6 tablets)	Suppresses symptoms of sneezing, runny nose and stuffy nose.

Additives: Hydroxypropyl cellulose, silicic anhydride, calcium hydrogen phosphate, iron sesquioxide, cellulose, magnesium stearate

Precautions for Storage and Handling

- (1) Store in a cool, dry place away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not replace the medicine into a different container. (This may result in misuse or a change in quality)
- (4) Do not take the product after the expiration date. Use this medicine within 6 months of opening, even if still within the expiration date. (In order to ensure quality)



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.