Make sure to read this instruction sheet before taking this medicine.
Additionally, store this sheet so that it can be referenced when needed.

For runny nose and stuffy nose caused by hay fever

Japanese Pharmacopoeia Fexofenadine hydrochloride tablets

PABRON RHINITIS FEXOFENADINE TABLETS

Brand name: PABRON RHINITIS FEXOFENADINE TABLETS Class 2 OTC drug <Allergy drug for rhinitis>

- **PABRON RHINITIS FEXOFENADINE** TABLETS are a rhinitis medication specifically for allergy sufferers, containing the second -generation antihistamine fexofenadine hydrochloride.
- Less likely to cause drowsiness or a decrease in concentration, judgment, or work efficiency.
- Take onetablet twicea dayfor all-day effectiveness.



Precautions for Use

Things You Must Not Do

(Failure to follow these precautions may lead to your current symptoms worsening or a greater chance of side effects or accidents occurring)

1 The following persons should not take this medicine:

(1) Persons who have had allergic symptoms as a result of taking this medicine or any component of it.

(2) Children under 15 years of age.

2 Do not use any of the following medicines while taking this medicine:

Other allergy medication (including medication for skin disorders and oral medication for rhinitis), oral medication containing antihistamines (including cold drugs, antitussive and expectorant combinations, motion sickness medication, hypnotic sedatives, etc.), formulations containing aluminum hydroxide or magnesium hydroxide (antacids), erythromycin (antibiotics), apalutamide (medication for prostate cancer treatment)

3 Do not drink alcohol before or after taking this medicine

4 Do not take this drug if you are breastfeeding, or avoid breastfeeding if you are taking this drug

(This medicine has been confirmed to have transference into breast milk in animal tests)



Things To Consult On

1 The following individuals should consult with their doctor, pharmacist, or a registered seller before taking this medicine

(1) Persons receiving treatment from a doctor.

(2) Persons who do not know whether their rhinitis is due to allergic rhinitis or other causes, such as a cold.

(3) Persons who have been diagnosed with other allergic diseases, such as bronchial asthma and atopic dermatitis.

(4) Persons with severe nasal congestion.

(5) Persons who are pregnant or who think they may be pregnant.

(6) Elderly persons.

(7) Persons with a history of allergic reactions caused by medicines, etc.

2 If any of the following symptoms appear after you have taken this medicine, you may be experiencing side effects. As such, discontinue use immediately and consult with your doctor, pharmacist or registered seller, bringing this instruction sheet with you

Relevant Area	Symptoms
Skin	Swelling of throat, eyelids or lips, etc., rashes, itching, hives, redness of skin
Digestive system	Nausea/vomiting, loss of appetite
Neuropsychiatric system	Numbness, headache, fatigue, lethargy, dizziness, insomnia, nervousness, nightmares, sleep disorders
Urinary tract	Frequent urination, difficulty urinating, nightmares, sleep disorders
Other	Palpitations, altered sense of taste, edemas, chest pain, difficulty breathing, elevated blood pressure, menstrual abnormalities

In rare cases, the following serious symptoms may occur.
In such cases, see a doctor immediately.

Symptom Name	Symptoms
Shock (anaphylaxis)	Itching, hives, hoarse voice, sneezing, itchy throat, breathlessness, palpitations, and confusion immediately after taking this medicine.
Liver dysfunction	Fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, general sluggishness, loss of appetite, etc.
Agranulocytosis, leukopenia, neutropenia	Sudden high fever, chills, sore throat, etc.

3 The following symptoms may appear after taking this medicine. If these symptoms persist or worsen, discontinue use and consult with your doctor, pharmacist or registered seller.
Dry mouth, constipation, diarrhea, drowsiness

Efficacy

Reduction of nasal allergy symptoms caused by pollen, house dust, etc.,: Sneezing, runny nose and stuffy nose

Usage/Dosage

Take the following dosages with room-temperature or lukewarm water.

Age	Single dose	Number of doses
Adults (15 years and older)	1 tablet 	Twice per day, morning and evening
Under 15 years of age	Do not take this medicine	

Note:
(1) Strictly follow the prescribed usage and dosage.
(2) When using the product for seasonal allergic rhinitis symptoms such as those caused by pollen, for best results, start taking the medicine on a day when airborne pollen is predicted or as soon as symptoms first appear.
(3) The product is most effective after continued use.
(4) If symptoms do not improve after using this medicine for 1 week, consult with your doctor, pharmacist or registered seller. Also consult with your doctor, pharmacist or registered seller if you plan to use the product for more than 2 weeks, even if your symptoms improve.
(5) How to take out the tablets:
As shown in the illustration, press firmly on the convex part of thePTPsheet containing the tablet with a fingertip to tear the aluminum foil on the reverse side, remove the tablet and take it. (Accidental ingestion of the PTP sheet as-is may lead to unexpected accidents, such as penetration into the esophageal mucosa.)

<Tablet removal diagram>


Ingredients


In 2 tablets

Fexofenadine hydrochloride 120 mg

Additives: Cellulose, partially pregelatinized starch, povidone,croscarmellose sodium, anhydrous silicic acid,magnesiumstearate, hypromellose, talc, titanium oxide, macrogol, yellow ferric oxide, ferric oxide

Precautions for Storage

(1) Store in a cool, dry place that is out of direct sunlight.
(2) Keep out of reach of children.
(3) Do not transfer to another container. (This may result in misuse or a change in quality)
(4) Do not use this product beyond its expiration date.



Disclaimer on Multilingual OTC Product Information

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.
- The provider (or Our Company) assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.